

Recombine often or perish: Genome evolution in bacterial and eukaryotic pathogens

Molecular Epidemiology of Infectious Diseases
Lecture 7

February 26th, 2024

**Classic population
genetic models of
evolution focus on
changes in genotype
frequencies**

Evolutionary Theory 101: Selection




Selection “acts” on genetic variation (i.e. mutations or alleles) at one or a few loci to increase the frequency of higher fitness variants.

Selection can purge deleterious mutations and increase the frequency of beneficial mutations.

The strength of selection will depend on the fitness effects of mutations and the size of populations due to genetic drift.

Genetic drift


Genetic drift refers to stochastic fluctuations in genotype frequencies caused by random variation in reproduction and survival. Stochastic variation and drift play a larger role in smaller populations.



Genetic drift


The probability that a beneficial mutation reaches fixation (freq \rightarrow 1.0) depends both on its selective advantage (s or σ) and the effective population size (N_e) – the number of individuals that contribute progeny to the next generation.

$$S = W_{mut} - W_{wt}$$



Evolutionary phase diagrams


Phase diagrams describe how the large-scale properties of systems depend on key variables and identify **critical points** at which the qualitative behaviour of the system changes.



Genetic drift


The probability that a beneficial mutation reaches fixation (freq \rightarrow 1.0) depends both on its selective advantage (s or σ) and the effective population size (N_e) – the number of individuals that contribute progeny to the next generation.

$$S = W_{mut} - W_{wt}$$




Selection vs. drift

The relative importance of selection versus drift is determined by $N_e s$



**Ok, but how do
pathogen genomes
actually evolve?**

Recombination vs. mutation



Recombination vs. mutation rates

The ratio r/m varies widely among different microbial pathogens


Table 1 The ratio of nucleotide changes as the result of recombination relative to point mutation (*r/m*) for different bacteria and archaea estimated from MLST data using ClonalFrame

Species	Phylum/division	Ecology	n STs	n loci	r/m	95% CI	Reference
<i>Flavobacterium psychrophilum</i>	Bacteroidetes	Obligate pathogen	33	7	63.6	32.8–82.8	Nicolas <i>et al.</i> (2008)
<i>Pelagibacter ubique</i> (SAR 11)	α -proteobacteria	Free-living, marine	9	8	63.1	47.6–81.8	Vergin <i>et al.</i> (2007)
<i>Vibrio parahaemolyticus</i>	γ -proteobacteria	Free-living, marine (OP)	20	7	39.8	27.4–48.2	Gonzalez-Escalona <i>et al.</i> (2008)
<i>Salmonella enterica</i>	γ -proteobacteria	Commensal (OP)	50	7	30.2	21.0–36.5	web.mpiib-berlin.mpg.de/mlst
<i>Vibrio vulnificus</i>	γ -proteobacteria	Free-living, marine (OP)	41	5	26.7	19.4–33.3	Bisharat <i>et al.</i> (2007)
<i>Streptococcus pneumoniae</i>	Firmicutes	Commensal (OP)	52	6	23.1	16.7–29.0	Hanage <i>et al.</i> (2005)
<i>Microcystis aeruginosa</i>	Cyanobacteria	Free-living, aquatic	79	7	18.3	13.7–21.2	Tanabe <i>et al.</i> (2007)
<i>Streptococcus pyogenes</i>	Firmicutes	Commensal (OP)	50	7	17.2	6.8–24.4	Enright <i>et al.</i> (2001)
<i>Helicobacter pylori</i>	α -proteobacteria	Commensal (OP)	117	8	13.6	12.2–15.5	pubmlst.org
<i>Moraxella catarrhalis</i>	γ -proteobacteria	Commensal (OP)	50	8	10.1	4.5–18.6	web.mpiib-berlin.mpg.de/mlst
<i>Neisseria meningitidis</i>	β -proteobacteria	Commensal (OP)	83	7	7.1	5.1–9.5	Jolley <i>et al.</i> (2005)
<i>Plesiomonas shigelloides</i>	γ -proteobacteria	Free-living, aquatic	58	5	7.1	3.8–13.0	Salerno <i>et al.</i> (2007)
<i>Neisseria lactamica</i>	β -proteobacteria	Commensal	180	7	6.2	4.9–7.4	pubmlst.net
<i>Mycococcus xanthus</i>	δ -proteobacteria	Free-living, terrestrial	57	5	5.5	1.9–11.3	Vos and Velicer (2008)
<i>Haemophilus influenzae</i>	γ -proteobacteria	Commensal (OP)	50	7	3.7	2.6–5.4	Meats <i>et al.</i> (2003)
<i>Wolbachia</i> b complex	α -proteobacteria	Endosymbiont	16	5	3.5	1.8–6.3	Baldo <i>et al.</i> (2006)
<i>Campylobacter insulaenigrae</i>	α -proteobacteria	Commensal (OP)	59	7	3.2	1.9–5.0	Stoddard <i>et al.</i> (2007)
<i>Mycoplasma hyopneumoniae</i>	Firmicutes	Commensal (OP)	33	7	3.0	1.1–5.8	Mayor <i>et al.</i> (2007)
<i>Haemophilus parasuis</i>	γ -proteobacteria	Commensal (OP)	79	7	2.7	2.1–3.6	Olvera <i>et al.</i> (2006)
<i>Campylobacter jejuni</i>	α -proteobacteria	Commensal (OP)	110	7	2.2	1.7–2.8	pubmlst.org
<i>Halorubrum</i> sp.	Halobacteria (Archaea)	Halophile	28	4	2.1	1.2–3.3	Papke <i>et al.</i> (2004)
<i>Pseudomonas viridiflava</i>	γ -proteobacteria	Free-living, plant pathogen	92	3	2.0	1.2–2.9	Goss <i>et al.</i> (2005)
<i>Bacillus weihenstephanensis</i>	Firmicutes	Free-living, terrestrial	36	6	2.0	1.3–2.8	Sorokin <i>et al.</i> (2006)
<i>Pseudomonas syringae</i>	γ -proteobacteria	Free-living, plant pathogen	95	4	1.5	1.1–2.0	Sarkar and Guttman (2004)
<i>Sulfolobus islandicus</i>	Thermoprotei (Archaea)	Thermoacidophile	17	5	1.2	0.1–4.5	Whitaker <i>et al.</i> (2005)
<i>Ralstonia solanacearum</i>	β -proteobacteria	Plant pathogen	58	7	1.1	0.7–1.6	Castillo and Greenberg (2007)
<i>Enterococcus faecium</i>	Firmicutes	Commensal (OP)	15	7	1.1	0.3–2.1	Homan <i>et al.</i> (2002)
<i>Mastigocladus laminosus</i>	Cyanobacteria	Thermophile	34	4	0.9	0.5–1.5	Miller <i>et al.</i> (2007)
<i>Legionella pneumophila</i>	γ -proteobacteria	Protozoa pathogen	30	2	0.9	0.2–1.9	Coscollo and Gonzalez-Candelas (2007)
<i>Microcoleus chthonoplastes</i>	Cyanobacteria	Free-living, marine	22	2	0.8	0.2–1.9	Lodders <i>et al.</i> (2005)
<i>Bacillus thuringiensis</i>	Firmicutes	Insect pathogen	22	6	0.8	0.4–1.3	Sorokin <i>et al.</i> (2006)
<i>Bacillus cereus</i>	Firmicutes	Free-living, terrestrial (OP)	13	6	0.7	0.2–1.6	Sorokin <i>et al.</i> (2006)
<i>Oenococcus oeni</i>	Firmicutes	Free-living, terrestrial	17	5	0.7	0.2–1.7	de Las Rivas <i>et al.</i> (2004)
<i>Escherichia coli</i> ET-1 group	γ -proteobacteria	Commensal (free-living?)	44	7	0.7	0.03–2.0	Walk <i>et al.</i> (2007)
<i>Listeria monocytogenes</i>	Firmicutes	Free-living, terrestrial (OP)	34	7	0.7	0.4–1.1	Salcedo <i>et al.</i> (2003)
<i>Enterococcus faecalis</i>	Firmicutes	Commensal (OP)	37	7	0.6	0.0–3.2	Ruiz-Garbajosa <i>et al.</i> (2006)
<i>Porphyromonas gingivalis</i>	Bacteroidetes	Obligate pathogen	99	7	0.4	0.0–3.4	Enersen <i>et al.</i> (2006)
<i>Yersinia pseudotuberculosis</i>	γ -proteobacteria	Obligate pathogen	43	7	0.3	0.0–1.1	web.mpiib-berlin.mpg.de/mlst
<i>Chlamydia trachomatis</i>	Chlamydiae	Obligate pathogen	14	7	0.3	0.0–1.8	Pannekoek <i>et al.</i> (2008)
<i>Klebsiella pneumoniae</i>	γ -proteobacteria	Free-living, terrestrial (OP)	45	7	0.3	0.0–2.1	Diancourt <i>et al.</i> (2005)
<i>Bordetella pertussis</i>	β -proteobacteria	Obligate pathogen	32	7	0.2	0.0–0.7	Diavatopoulos <i>et al.</i> (2005)
<i>Brachyspira</i> sp.	Spirochaetes	Commensal (OP)	36	7	0.2	0.1–0.4	Rasback <i>et al.</i> (2007)
<i>Clostridium difficile</i>	Firmicutes	Commensal (OP)	34	6	0.2	0.0–0.5	Lenne <i>et al.</i> (2004)
<i>Bartonella henselae</i>	α -proteobacteria	Obligate pathogen	14	7	0.1	0.0–0.7	Arvand <i>et al.</i> (2007)
<i>Lactobacillus casei</i>	Firmicutes	Commensal	32	7	0.1	0.0–0.5	Diancourt <i>et al.</i> (2007)
<i>Staphylococcus aureus</i>	Firmicutes	Commensal (OP)	53	7	0.1	0.0–0.6	Enright <i>et al.</i> (2000)
<i>Rhizobium gallicum</i>	α -proteobacteria	Free-living, terrestrial	33	3	0.1	0.0–0.3	Silva <i>et al.</i> (2005)
<i>Leptospira interrogans</i>	Spirochaetes	Commensal (OP)	61	7	0.02	0.0–0.1	Thaipadungpanit <i>et al.</i> (2007)

Vos & Didelot (ISME, 2008)

Recombination vs. mutation rates

The relative ratio of recombination versus mutation rates determines whether selection acts primarily on individual mutations or whole genotypes/haplotypes.




Selection on linked sites

When $r < \mu$ mutations will be linked to other mutations in the same genetic background (haplotype). The fate of a mutation therefore depends on the fitness effects of other linked mutations (background or interference selection).

In contrast, when $r > \mu$ recombination will reshuffle mutations onto different genetic backgrounds such that selection can act on individual mutations (allelic selection).

Recombination vs. mutation rates

The relative ratio of recombination versus mutation rates determines whether selection acts primarily on individual mutations or whole genotypes/haplotypes.



Clonal vs. horizontal evolution


Clonal evolution occurs by vertical descent where genetic material is passed from parents to children.

In contrast, genetic material can be exchanged **horizontally** between lineages through recombination or horizontal gene transfers.

Clonal interference

Clonal interference arises in large asexual populations with high mutations rates and large population sizes.

Multiple lineages with beneficial mutations compete with one another.



Cvijovic et al. (Trends in Genetics, 2018)


Clonal interference through competition

The Microbial Evolution and Growth Arena (MEGA) experiments show multiple strains of antibiotic resistant *E. coli* competing with one other for space.



Clonal interference in *M. tuberculosis*

Dynamics of TB clones within a host treated with a series of antibiotics. Only 7 out of 12 beneficial DRM mutants reach fixation while the rest are outcompeted.



Clonal interference: summary

Role of genetic drift becomes negligible as competition creates strong selection for highly fit genotypes.

Increases chance that “best” genotype with the largest fitness advantage goes to fixation. This genotype may often carry multiple beneficial mutations.

However, interference can actually slow down the rate of adaptation (increases in mean pop fitness) as multiple beneficial genotypes will compete against one another.

**How do multi-drug
resistant bacteria
evolve?**

Resistance evolution in MRSA

Methicillin resistant *Staphylococcus aureus* first emerged in the 1960's and causes dangerous bloodstream infections with roughly 30% mortality.


S. aureus is highly clonal with a low recombination rate and only a few lineages are responsible for most infections.

Methicillin resistance is acquired via the staphylococcal cassette chromosome *SCCmec*, a mobile genetic element that integrates a cassette of genes into the bacterial chromosome (most likely by transduction from bacteriophages).

SCCmec acquisition is a classic example of horizontal gene transfer (HGT).

Acquisition of methicillin resistance

S. aureus phylogenies show independent acquisitions of *SCCmec* cassettes in across different lineages suggesting frequent horizontal transfers.




The pangenome concept

The **core genome** contains the highly conserved genes with often essential functions (i.e. DNA replication) that are present within all members of a species.

The **accessory genome** includes gene content that varies between strains including chromosomal cassettes, prophages, transposons and pathogenicity islands. These genes can be chromosomal or extrachromosomal (e.g. on plasmids).

Prokaryote populations are often characterized by extensive sharing of accessory genes due to horizontal gene transfers and subsequent gene losses.

The MRSA accessory genome



Evolution of multidrug resistant MRSA

Increased fitness of initially methicillin resistance strains leads to large clonal expansions.


Accumulation of additional genes conferring resistance to other antibiotics occurred through further HGTs, leading to superfit multidrug resistance clones.

These multi-drug resistant (MDR) strains have acquired resistance to several major classes of antibiotics including fluoroquinolones and erythromycin.

MDR MRSA

“MRSA epitomizes a now all-too-familiar evolutionary route by which successful AMR clones emerge in response to local antimicrobial usage, undergo population expansion under selection from sustained antimicrobial exposure and then explode into pandemic spread”

Baker et al. (Science, 2017)



Recombine often or perish

Moderately high mutation rates but low recombination rates cause competition and clonal interference between high-fitness lineages.

Rapid adaptation to selective pressures like antimicrobials occurs by frequent horizontal transfers of beneficial genetic elements in the accessory genome.

Successful lineages therefore tend to be clones that have acquired multiple beneficial genes or other genetic elements through HGT.


“Two-speed” genomes in eukaryotes

Pathogenic fungi often have virulence-related and effector genes involved in host-adaptation compartmentalized to genomic regions with elevated mutations rates and high densities of mobile genetic elements, including:

Accessory chromosomes: lineage-specific chromosomes with low gene density but high mutation rates that can move horizontally between lineages.

TE-rich compartments: regions of the core genome that are gene-sparse but highly variable due to transposable elements or other mobile genetic elements.

Fast evolving genome compartments



**How clonal are
bacteria and other
microbial pathogens?**

Clonal vs. horizontal evolution

Clonal evolution occurs by vertical descent where genetic material is passed from parents to children.

In contrast, genetic material can be exchanged **horizontally** between lineages through recombination or horizontal gene transfers.


What determines clonality?

Rates of recombination and horizontal exchange have traditionally been thought of as the key determinants of clonality.

However, recombination also interacts with selection to determine how clonal a population is at any particular point in time.



Selective sweeps eliminate diversity

Strong selection can lead to rapid clonal expansions and genome-wide selective sweeps of linked variants.



Clonal expansions of chestnut blight

The invasive S12 genotype is an asexual and more virulent lineage of *Cryphonectria parasitica* undergoing a clonal expansion in European chestnut trees.




Recombination vs. selection

The strength of selection relative to the recombination rate determines how selective sweeps impact diversity elsewhere in the genome.


*Genome-wide
selective sweeps*

$$s = W_{mut} - W_{wt}$$



*Gene-specific
selective sweeps*

Gene-specific vs. genome-wide sweeps



Recommended reading



ELSEVIER

Available online at www.sciencedirect.com

ScienceDirect

Current Opinion in
Microbiology

How clonal are bacteria over time?

B Jesse Shapiro



Bacteria and archaea reproduce clonally (vertical descent), but exchange genes by recombination (horizontal transfer). Recombination allows adaptive mutations or genes to spread rapidly within (or even between) species, and reduces the burden of deleterious mutations. Clonality — defined here as the balance between vertical and horizontal inheritance — is therefore a key microbial trait, determining how quickly a population can adapt and the size of its gene pool. Here, I discuss whether clonality varies over time and if it can be considered a stable trait of a given population. I show that, in some cases, clonality is clearly not static. For example, non-clonal (highly recombining) populations can give rise to clonal expansions, often of pathogens. However, an analysis of time-course metagenomic data from a lake suggests that a bacterial population's past clonality (as measured by its genetic diversity) is a good predictor of its future clonality. Clonality therefore appears to be relatively — but not completely — stable over evolutionary time.

where a bacterial population of interest happens to fall along a spectrum of clonality can help us understand its biology, and even make predictions about its evolution.

The opposite of clonality is panmixis — a situation in which the rate of horizontal transfer is higher than the rate of vertical cell division, resulting in random association (linkage equilibrium) among loci in the genome [1,2]. However, rates of horizontal transfer (recombination) vary widely across the genome, such that a population can be mostly clonal, except for a few loci in the genome [3]. These loci came to be termed genomic islands — a metaphor I will build upon below. Some of the first islands identified were called pathogenicity islands because they contained virulence factors [4]. However, non-pathogenic environmental bacteria also contain islands, conferring adaptation to different ecological niches. For example, genes in *Prochlorococcus* genomic islands confer

The extended island metaphor

Clonality varies considerably between different species and may reflect their recent demographic history.

Extended island metaphors of microbial genome evolution				
Geographic metaphor	Genetic unit to which the metaphor applies	Type of selective sweep experience by the unit	Dominant mode of genetic transmission	Example
Island	Gene	Gene-specific	Horizontal	Genes in the <i>V. cholerae</i> integron [22*,23]
Peninsula	Gene	Genome-wide	Vertical (clonal)	The cholera toxin gene, acquired horizontally, then linked to a clonal <i>V. cholerae</i> genome [9,21]
Continent	Genome	Genome-wide	Vertical (clonal)	Clonal expansions of <i>S. aureus</i> [28], <i>M. tuberculosis</i> [31,43]
Archipelago	Genome	Gene-specific	Horizontal	Hotspring cyanobacteria [11*], ocean vibrios [13*,14], pneumococcus [44,46*]

Clonality can dynamically vary over time

Genome-wide selective sweeps increase clonality even if recombination rates were historically high.

Genomic islands, pieces of DNA that were transferred horizontally, can therefore become peninsulas linked by the conserved regions of the genome (i.e. the continents).

Continents can be broken up over time into archipelagos by recombination.

